

Autocorrelation in Updating Pure SU(3) Lattice Gauge Theory by the use of Overrelaxed Algorithms

QCD_TARO Collaboration

K.Akemi ^a, Ph.deForcrand ^b, M.Fujisaki^a, T.Hashimoto ^c, H.C.Hege ^d, S.Hioki ^e, O.Miyamura^e, A.Nakamura ^f, M.Okuda^a, I.O.Stamatescu ^g, Y.Tago^a and T.Takaishi^e

^aComputational Science Research Laboratory, Fujitsu Limited, Ota-ku, Tokyo 144, Japan

^bIPS,ETH-Zürich, CH-8092 Zürich, Switzerland

^cDept. of Applied Physics, Fukui University, Fukui 910, Japan

^dZIB, D-1000 Berlin 31, Germany

^eDept. of Physics, Hiroshima University, Higashi-Hiroshima 724, Japan

^fDept. of Physics, Waseda University, Tokyo 169, Japan

^gFEST Heidelberg and Institut für Theoretische Physik, Universität Heidelberg, D-6900 Heidelberg, Germany

We measure the sweep-to-sweep autocorrelations of blocked loops below and above the deconfinement transition for SU(3) on a 16^4 lattice using 20000-140000 Monte-Carlo updating sweeps. A divergence of the autocorrelation time toward the critical β is seen at high blocking levels. The peak is near $\beta = 6.33$ where we observe 440 ± 210 for the autocorrelation time of 1×1 Wilson loop on 2^4 blocked lattice. The mixing of 7 Brown-Woch overrelaxation steps followed by one pseudo-heat-bath step appears optimal to reduce the autocorrelation time below the critical β . Above the critical β , however, no clear difference between these two algorithms can be seen and the system decorrelates rather fast.

1. Introduction

In the Monte Carlo evaluation of physical observables, we should gather independent samples to get statistically meaningful results. We generate a sequence of configurations and then we choose samples separated by some sweeps to ensure their independence. This separation should be larger than the Monte Carlo autocorrelation time τ . Otherwise statistical errors should be corrected to take correlations into account. It is crucial in lattice QCD to estimate correctly these errors and the underlying autocorrelation time.

In this paper we report autocorrelation measurements in pure SU(3) lattice gauge theory.[1]

The aims of our study are

1. to extract autocorrelation time
2. to see operator dependence of autocorrelation,i.e. blocked loops
3. to see β dependence of autocorrelation which tells us some information about the critical slowing down toward the continuum limit.




Fig.1 MC history of W11 at level=2-4, $\beta=6.38$.

2. Overrelaxation, Blocking scheme and Numerical simulation

To update the gauge field here we use Cabbibo-Marinari[2] pseudo-heat-bath(PHB) with 3 SU(2) subgroups and to perform the overrelaxation(OR) we adopt Brown-Woch[3] microcanonical method with 2 randomly chosen SU(2) subgroups.

We introduce a parameter K which denotes the mixing ratio between OR and PHB, namely we choose OR or PHB stochastically with the probabilities $K/(K+1)$ and $1/(K+1)$.

For blocking we use the Swendsen[4] factor 2 blocking scheme. For details and results of our MCRG study, see Ref.[5]

We work on a 16^4 lattice and 20000-140000 sweeps are performed at all β . All data are $K=9$ case except for $\beta=6.3$ and 6.8 . At $\beta=6.3$ and 6.8 , $K=0, 3, 5, 7, 9, 12$ were studied to check the efficiency of OR algorithm.

After a few thousands of sweeps for thermalization, we measure operators: W11, W12, W22, Wchair, Wsofa and Wtwist at every blocking level from 1 to 4 (i.e. from 16^4 to 2^4). Blocking is performed every 10 sweeps. All our simulations have been performed on 64-node AP1000 at Fujitsu.

3. Results

In Fig.1, we show typical Monte Carlo history of W11(1×1 wilson loop) at blocking levels 2-4 at $\beta=6.38$.




Fig.2 $\rho(t)$ vs. t , same data as in Fig.1.

In this case no clear signal of sweep-to-sweep correlation at low blocking level can be seen whereas at high blocking level some structure emerges which we cannot see at lower levels.

Then we define the autocorrelation function $\rho(t)$ of observable O as

$$\rho(t) = \frac{\langle (O(i) - O_A)(O(i+t) - O_B) \rangle}{\sqrt{\langle (O(i) - O_A)^2 \rangle \langle (O(i+t) - O_B)^2 \rangle}} \quad (1)$$

where $\langle \rangle$ denotes average over i and $O_A = \langle O(i) \rangle$ and $O_B = \langle O(i+t) \rangle$.

Fig.2 shows the autocorrelation function $\rho(t)$ for $t=0-1000$ sweeps for the same data as in Fig.1. In Fig.2 we see long range autocorrelation, as stated above, at high blocking level.

This tendency is very natural. Since W11 at high blocking level has effectively large area, then more Monte Carlo sweeps are needed to alter its value. In other words, W11 at high blocking level is not easily influenced by short range fluctuations whereas at lower level it is.

Now we introduce integrated autocorrelation time τ_{int} defined as:

$$\tau_{int} = \rho(0) + 2 \sum_{t=1}^N \rho(t) \frac{N-t}{N} \quad (2)$$

In this analysis we choose $N=1000$, so that τ_{int} can extract long range information. For the error estimation of correlated samples, τ_{int} is the relevant quantity and enters in our final error analysis.




Fig.3(a) $\rho(t)$ vs. t , for $K=0,3,7,9$ at $\beta=6.3$.

We have done this analysis not only for W11 but also for other measured loops such as W12, W22, Wsofa etc. There exist, of course, differences among them about autocorrelation function and time, however, we found that for a given blocking level, these differences are not significant. So from now on, we concentrate on W11 at level=4.

As stated above, we have checked the efficiency of OR algorithm changing the mixing parameter K .

In Fig.3(a), $\rho(t)$ for $t=0-600$ at $\beta=6.3$ for $K=0,3,7$ and 9 are displayed. In this figure, $K=7$ seems to work well to reduce the autocorrelation. To see this quantitatively, we show τ_{int} defined by eq.(2) for several values of K at $\beta=6.3$ in Fig.3(b). We can see the efficiency of $K=7$ case also in this figure.

From this analysis, we can extract the typical autocorrelation time of W11 at level=4 at $\beta=6.3$ such as: about 300 sweeps for PHB, about 100 sweeps for OR at $K=7$. In ref.[6] Decker and de Forcrand give similar numbers. These numbers are somewhat larger than those obtained by other authors. This is because we are using blocked loops as observables. To see the long range autocorrelation, extended operators look suitable because small operators are easily influenced by fluctuations.

When we want to include error bars, we must be careful to control these long range autocorre-




Fig.3(b) τ_{int} for several K values at $\beta=6.3$.

lation effects.

Let $\delta\rho_0(t)$ be the error on $\rho(t)$ calculated as if samples were independent. Then if we consider the dependence between samples, the error is modified as

$$(\delta\rho(t))^2 = (\delta\rho_0(t))^2 \tau_{int}. \quad (3)$$

This error on $\rho(t)$ generates an error on the extracted autocorrelation times. We estimate it by

$$\delta\tau_{int} = \frac{\delta\rho(\tau_{int})}{\rho(\tau_{int})} \tau_{int}. \quad (4)$$

To see the definition dependence of error analysis, we also evaluate errors using jackknife method. To perform jackknife analysis for error estimates of autocorrelation function, we first divide the sample into bins. We get $\delta\rho(t)$ and then we can estimate the error of autocorrelation time following eq.(4). It is thought that if we double the size of a bin, error increases roughly by a factor $\sqrt{2}$ when size of a bin is smaller than the autocorrelation time and then levels off when size of a bin exceeds the autocorrelation time. In our analysis there are many cases when the error won't level off. In such cases the error on the error of autocorrelation function is very large.

We show the results of this jackknife study also in Fig.3(b). The errorbar with full circle comes from using eq.(3) and that with open circle comes from jackknife analysis. In all cases of K , jack-




Fig.4(a) $\rho(t)$ vs. t , for several β at $K=9$.

knife method estimates smaller errorbars than eq.(3).

If we include the error and consider again the efficiency of OR algorithm, we can only say that $K=7$ looks optimal. To adopt OR algorithm, it is important to choose the optimal value of K to update the system efficiently. The optimum range of K is surprisingly narrow.

We have performed the same analysis also at $\beta=6.8$. At this β , τ_{int} is rather small for any K (typically less than 50 sweeps) and no clear advantage of OR can be seen.

Finally we consider the β dependence of $\rho(t)$ and τ_{int} .

We show the behaviors of $\rho(t)$ for different values of β in Fig.4(a). We can see long range autocorrelations at $\beta=6.33$ and 6.43 , whereas rather short range ones at $\beta=6.10$ and 6.80 .

The autocorrelation time as a function of β is displayed in Fig.4(b). τ_{int} seems to diverge around some β , associated with the deconfinement transition. This indicates that around this β_τ we must be careful to evaluate the statistical average of physical observables as we stressed in the beginning of this paper.

From these results, it is clear that if we want to stay in the confinement regime, we must stay below $\beta_\tau = 6.34 \pm 0.03$ on a 16^4 lattice.

Combining with renormalization group analysis at $\beta=6.8$ on 32^4 lattice indicating $\Delta\beta = 0.53 \pm .02$ (see Ref.5), we must stay at $\beta \leq 6.87 \pm 0.05$




Fig.4(b) τ_{int} as a function of β at $K=9$.

on a 32^4 lattice.

ACKNOWLEDGEMENTS

We are indebted to M.Ikesaka, Y.Inada, K.Inoue, M.Ishii, T.Saito, T.Shimizu and H.Shiraishi at the Fujitsu parallel computing research facilities for their valuable comments on parallel computing.

REFERENCES

1. R.Gupta et al., Mod.Phys.Lett.A 3 (1988) 1367.
2. N.Cabibbo and E.Marinari, Phys.Lett. 119B (1982) 387.
3. F.R.Brown and T.J.Woch, Phys.Rev.Lett. 58 (1987) 2394.
4. R.Swendsen, Phys.Rev.Lett. 47 (1981) 1775.
5. K.Akemi et al., "Monte Carlo Renormalization Group Study at Large β in Confinement Region", these proceedings.
6. K.M.Decker and Ph. de Forcrand, Nucl. Phys. B (Proc. Suppl.) 17 (1990) 567.